

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

### **Supplemental Material**

#### **Prenatal Exposure to Perfluoroalkyl Substances and Child Behavior at Age 12: A PELAGIE Mother–Child Cohort Study**

Hélène Tillaut, Christine Monfort, Florence Rouget, Fabienne Pelé, Fabrice Lainé, Eric Gaudreau, Sylvaine Cordier, Charline Warembourg, Dave Saint-Amour, and Cécile Chevrier

#### **Table of Contents**

**Table S1.** PELAGIE mother-child cohort, comparisons between included and not included in the study.

**Table S2.** Parent-reported Strengths and Difficulties Questionnaire (SDQ) - Scores at age 12, PELAGIE mother-child cohort, Brittany, France (n=444). Values reported are the median (minimum-maximum) scores.

**Table S3.** Self-reported Dominic Interactive for Adolescents (DIA) - Scores at age 12, PELAGIE mother-child cohort, Brittany, France (n=444). Values reported are the median (minimum-maximum) scores.

**Table S4.** Parent-reported Strengths and Difficulties Questionnaire (SDQ) - Scores at age 12 – alpha de Cronbach.

**Table S5.** Self-reported Dominic Interactive for Adolescents (DIA) - Scores at age 12 – alpha de Cronbach.

**Table S6.** Associations between prenatal exposure measured in cord serum and behavior scores at age 12 – Overall and stratified on child sex –PELAGIE mother-child cohort.

**Table S7.** Associations between prenatal exposure measured in cord serum and behavior scores at age 12 – sensitivity analysis – Breastfeeding, PELAGIE mother-child cohort.

**Table S8.** Associations between prenatal exposure measured in cord serum and behavior scores at age 12 – sensitivity analysis – parental mental health (SF36 Emotional well-being), PELAGIE mother-child cohort.

**Table S9.** Distribution of cord-blood serum concentrations of persistent organic pollutant.

**Table S10.** Associations between prenatal exposure measured in cord serum and behavior scores at age 12 – Results from the single pollutant approach, PELAGIE mother-child cohort.

**Table S11.** Associations between prenatal exposure measured in cord serum and behavior scores at age 12 – sensitivity analysis – co-exposure to persistent organic pollutant, PELAGIE mother-child cohort.

**Table S12.** Posterior inclusion probability – Bayesian Kernel Machine Regression (BKMR) models (n= 378) - PELAGIE mother-child cohort, Brittany, France.

**Table S13.** Overall PFAS mixture effect for SDQ Externalizing and DIA Internalizing scores – change in estimate (and 95% credible intervals) for concurrent increase of exposures compared to the 25th percentile and posterior inclusion probabilities (PIP) – Bayesian Kernel Machine Regression (BKMR) models (n= 378) - PELAGIE mother-child cohort, Brittany, France.

**Table S14.** Change in estimate (and 95% credible intervals) with all other exposures fixed at specified quantiles from the multipollutant BKMR analyses for SDQ Externalizing subscale (A) and DIA Internalizing subscale (B) – Bayesian Kernel Machine Regression (BKMR) models (n= 378) – PELAGIE mother-child cohort, Brittany, France.

**Table S15.** Exposure levels (wet weight, log2 transformed, centered on the median and IQR scaled) for compounds included in BKMR models (n=378) – PELAGIE mother-child cohort, Brittany, France.

**Figure S1.** Flow chart.

**Figure S2.** Directed acyclic graph.

**Figure S3.** Spearman correlation coefficients between detected values of cord-serum concentration of POPs and PFASs.

**Figure S4.** Associations between prenatal exposure to PFAS and child behavior – Results for the generalized additive models including restricted cubic splines.

**Figure S5.** Associations between prenatal exposure to PFAS and child behavior – Results for negative binomial regression models including an interaction term between exposure and sex – Scores for males and females and 95%CI.

**Figure S6.** Associations between prenatal exposure to PFAS measured in cord serum and behavior scores at age 12, overall and for females and males separately – Results from the single pollutant approach, PELAGIE mother-child cohort.

**Figure S7.** Change in estimate (and 95% credible intervals) with all other exposures fixed at specified quantiles from the multipollutant BKMR analyses for SDQ Externalizing subscale (A) and DIA Internalizing subscale (B) – Bayesian Kernel Machine Regression (BKMR) models (n= 378) – PELAGIE mother-child cohort, Brittany, France.

**Figure S8.** Exposure-response relationships for prenatal exposure to a mixture of PFAS and SDQ externalizing disorders score among 12-years-old children – Bayesian Kernel Machine Regression (BKMR) models (n= 378) - PELAGIE mother-child cohort, Brittany, France.

**Figure S9.** Exposure-response relationships for prenatal exposure to a mixture of PFAS and DIA internalizing subscale among 12-years-old children – Bayesian Kernel Machine Regression (BKMR) models (n= 378) - PELAGIE mother-child cohort, Brittany, France.